

HOMEWORK B

Instructor: Yoonseok Lee

Due: February 15, 2018

1. CH26 Q-11.

Pay attention to the dependence on the distance. For a point charge $E \propto 1/r^2$, for a dipole $E \propto 1/r^3$, and for a quadrupole $E \propto 1/r^4$. From the dimensional point of view, does it make sense?

2. CH26 E-19.

3. Using the online simulation at <http://www.falstad.com/vector3de/>, simulate the electric field lines of a quadrupole moment as shown in Figure 26-27 (Q-11). Sketch the field lines on the $x - y$ plain (z -slice in the simulation) with the minimal charge spacing. This configuration will show the quadrupole field lines at a large distance. Compare the direction of the electric field along the diagonal direction and the x -direction.

4. CH26 E-39.

5. CH26 P-2.

For $d \ll \sqrt{x^2 + z^2}$,

$$\frac{1}{\{x^2 + (z \pm d/2)^2\}^{3/2}} \approx \frac{1}{(x^2 + z^2)^{3/2}} \left(1 \mp \frac{3}{2} \frac{zd}{x^2 + z^2} \right).$$

Expand the denominator, then immediately $x^2 + z^2 \pm zd + (d/2)^2 \approx (x^2 + z^2) \pm zd$ (why?). Factor out $(x^2 + z^2)$ then $zd/(x^2 + z^2)$ becomes the small parameter for expansion.

6. CH26 P-11.

(i) What is the net force (Coulomb + gravitational force) on the electron? (ii) Then you can calculate the y -component velocity from the effective acceleration at the height of d and how long it would take to the height. (iii) During the time will it travel along the x -direction far enough to escape or not?

7. Ch26 P-12.

At this point you know you have to do Taylor expansion to get a form $F \propto z$ for $z \ll R$ near the center of the ring. This is the opposite limit of $z \gg R$.

8. CH26 P-14.

You do not have to solve this problem. But I ask you to think about this. I will discuss this in class. For a mass loaded spring, the equation of motion is $m\ddot{x} = -kx$ and the linear oscillation frequency $f = \frac{1}{2\pi} \sqrt{\frac{k}{m}}$. For a rotational oscillation, the equation of motion is based on the torque $\vec{\tau} = \vec{r} \times \vec{F}$: $\vec{\tau} = I\ddot{\theta} = -\alpha\theta$, and the rotational frequency is $f = \frac{1}{2\pi} \sqrt{\frac{\alpha}{I}}$ where I is the moment of inertia and α is the torsional spring constant. Check the physical dimensions of α and I if they produce $[f] = T^{-1}$.

9. CH27 E-3.

This figure may be helpful.




FIG. 1:

10. CH27 E-25.

11. CH27 P-14.

\vec{r} is the radial vector to a point inside the hole (P). Then $\vec{a} = \vec{r} - \vec{b}$ where \vec{b} is the vector to the point P from the center of the hole (O').




FIG. 2:

12. CH27 P-19.

Inside an isolated conductor $\vec{E} = 0$.