

**PHY 6646 - Quantum Mechanics II - Spring 2018**  
**Homework #14, due April 25**

1. The four lowest energy states of the hydrogen atom are split in energy by the interaction between electron spin and proton spin. See problem 15.1.2. The groundstate has total spin  $s = 0$ . At an energy  $\hbar\omega_0$ , where  $\omega_0 = (2\pi) 1.42$  GHz, above the ground state are three states with  $s = 1$ . The electromagnetic radiation emitted or absorbed by transitions between the  $s = 0$  and  $s = 1$  states is commonly referred to as the “21 cm line”

a) A plane electromagnetic wave of wavevector  $\vec{k}$ , frequency  $\omega = ck$ , and energy per unit surface and unit time  $I$ , is incident upon a hydrogen atom in its ground state. The electromagnetic wave causes a time-dependent perturbation of the hydrogen atom

$$H^1(t) = -\gamma_e \vec{S}_e \cdot \vec{B}(\vec{0}, t) \quad (0.1)$$

where  $\vec{S}_e$  is the spin of the electron,  $\gamma_e$  the electron gyromagnetic ratio, and  $\vec{B}(\vec{0}, t)$  the magnetic field at the location of the atom. Eq. (0.1) neglects the much weaker interaction ( $\gamma_p \ll \gamma_e$ ) of the electromagnetic field with the proton spin. Obtain the rate of transition of the atom from its  $s = 0$  ground state to any of the three  $s = 1$  states as a result of this perturbation. Does the rate depend on the direction of  $\vec{k}$ ? Does it depend on the state of polarization of the electromagnetic wave?

b) The atom in its ground state is placed in a thermal bath of electromagnetic radiation at temperature  $T$ . Show that the rate at which it makes transitions to the  $s = 1$  states is given by

$$R = \left( \frac{e}{m_e c} \right)^2 \frac{\hbar \omega_0^3}{c^3} \frac{1}{e^{\hbar\omega_0/k_B T} - 1} \quad (0.2)$$

where  $m_e$  is the electron mass, and  $k_B$  is Boltzmann's constant.

c) Verify that Eq. 0.2 is dimensionally correct.

2. Problem 20.1.1

3. Problem 20.2.1

4. Problem 20.2.2