PHY 7097 | Modern Astrophysics

Lecture XV.

Accretion



UF | UNIVERSITY of FLORIDA

Imre Bartos | Spring 2018

Disk Accretion

BH:



Radiation efficiency: U/m

Schwarzschild BH:	5.7%
Kerr BH:	≤ 42%

Most radiation will come from the innermost parts of the disk.

(Nuclear fusion: 0.7%)

(Gasoline: 0.00000003%)

Efficiency for WDs and NSs





WD or NS: particle collide with the stellar surface → kinetic energy is radiated away → order of magnitude of total luminosity is independent of the nature of the flow (spherical or disk).

Total luminosity: $L = G\dot{M}M/R$

X-ray binaries

- Roche lobe overflow
- Stellar wind

X-ray radiation is strong and detectable (no background)

- Low-mass X-ray binary
  - Smaller companion (e.g. main sequence)
  - Roche lobe
  - Faint in optical
- High-mass X-ray binary
  - Massive star companion (e.g. blue supergiant)
  - Massive star bright in optical
- Microquasars
  - Also emits radio – jet is important
  - Small relative of quasars



X-ray binaries



Thin disk accretion



Thin disk: $h \ll r$

- Simplifies calculations
- Allows efficient radiation

Viscosity in disk:

- Poorly understood
- Small-scale turbulence
- Magnetic fields

Shakura & Sunyaev (1973)

Assume viscosity is due to turbulence.
Viscosity coefficient in turbulence:

$$\nu = v_{\text{turb}} l_{\text{turb}}$$

$$v_{\text{turb}} \leq c_s \qquad l_{\text{turb}} \leq h$$

$$\nu = \alpha c_s h$$

$$0 \leq \alpha \leq 1 \quad \alpha\text{-disks}$$

α is a free parameter that is measured empirically. It falls within 0.1 – 1.

Thin disk accretion

$$T_s = (2 \times 10^9\ \mathrm{K})\alpha^{2/9}\left(\frac{M}{M_\odot}\right)^{-10/9} \dot{M}_{17}^{8/9}\left(\frac{r}{r_s}\right)^{-17/9} \mathcal{G}^{8/9}$$



Different regions with different emission mechanisms.

Heavier BH → softer spectrum.

Active galactic nuclei

Bright galactic center, due to an accreting supermassive BH.

Galaxy hosting an AGN is called an active galaxy.

Properties depend on:

- Mass of BH
- Accretion rate
- Orientation of disk
- Obscuration (e.g. dust)
- Presence or absence of jet.

Subcategories:

- Quasars
- Blazars



(Wikipedia)

Active galactic nuclei



Active galactic nuclei

A variety of classifications exist.

Radio-loud AGN --- the jet dominates the radio and possibly all other emission.

Radio-quiet AGN --- no jet or not relevant (e.g. pointing in some other direction).

Features of different types of galaxies

Galaxy type	Active nuclei	Emission lines		X-rays	Excess of		Strong radio	Jets	Variable	Radio loud
		Narrow	Broad		UV	Far-IR				
Normal	no	weak	no	weak	no	no	no	no	no	no
LINER	unknown	weak	weak	weak	no	no	no	no	no	no
Seyfert I	yes	yes	yes	some	some	yes	few	no	yes	no
Seyfert II	yes	yes	no	some	some	yes	few	no	yes	no
Quasar	yes	yes	yes	some	yes	yes	some	some	yes	some
Blazar	yes	no	some	yes	yes	no	yes	yes	yes	yes
BL Lac	yes	no	no/faint	yes	yes	no	yes	yes	yes	yes
OVV	yes	no	stronger than BL Lac	yes	yes	no	yes	yes	yes	yes
Radio galaxy	yes	some	some	some	some	yes	yes	yes	yes	yes

Classification

This is not certain.

Quasar (quasi stellar object) – high-power AGN.

Blazar (BL Lac object + quasar) – jetted AGN where we look down the jet.



Beckmann & Shrader 2012

Fermi bubbles



Right now the Milky Way is not an active galaxy.



Quasar variability

- Red noise

  (damped random walk)
- SMBH binary?



Graham+ Nature 2015

Tidal disruption events





http://physicstoday.scitation.org/doi/10.1063/PT.3.2382

Tidal disruption events



http://physicstoday.scitation.org/doi/10.1063/PT.3.2382

G2 gas cloud



G2 gas cloud